



100% active oxytetracycline hydrochloride

- Ministry Approved
- Effective against bacteria:
Aeromonas spp, Yersinia spp,
Vibrio spp and Myxobacteria spp
- 100% active oxytetracycline
- Use in Atlantic Salmon and
Rainbow trout
- Administration in fish food
- 720° days withdrawal time

Legal Category

POM-V

Package Quantities

1 kg, 2.5 kg plastic tubs

Marketing Authorisation Number

Vm 11003/4002

Marketing Authorisation Holder

PHARMAQ Ltd.

Dosage and Administration

Aquatet is given by oral administration at the rate of 75 mg/kg bodyweight of fish daily for initially 4 days. After the initial treatment, mortalities should be assessed and a further 4 day treatment course should be administered if necessary.

Aquatet is for administration only through the feed by mixing with manufactured feed prior to feeding. Feeding rates will vary according to water temperature and it may therefore be more convenient to medicate on a basis of a fixed rate e.g. 1% of bodyweight, with an extra daily feed requirement being met by unmedicated food. The following inclusions rates will provide the recommended dose:

Daily feed rate % bodyweight	Aquatet inclusion rate		
	per 5 kg of food	per 25 kg of food	per 1 tonne of food
0.5	75 g	375 g	15 kg
1	37.5 g	187.5 g	7.5 kg
2	18.75 g	93.75 g	3.75 kg

Method of mixing:

Weigh out appropriate amounts of fish pellets and Aquatet powder and mix well together in the dry state. As an aid to the adhesion of the Aquatet powder in the fish pellets, a small quantity of tepid gelatin solution or edible oil is then added to the food while mixing. Sufficient quantity should be mixed into the medicated food until it is slightly dampened.

Contra-indications, Warnings etc.

Animals must not be slaughtered for human consumption during treatment. Fish may be slaughtered for human consumption 720° days after the end of treatment.

When mixing Oxytetracycline, rubber gloves should be worn to avoid the possibility of skin reactions and a respirator which at least complies with European Standard EN149, FFP2 to avoid inhalation of any dust particles.

Pharmaceutical Precautions

Store the original container at a temperature not exceeding 25°C and in a dry place. Re-close the part-used containers. Protect from light.

Medicated diet should be used for the treatment period only. Keep out of the reach of children.

Licensed in

United Kingdom and Ireland.

Only available through your veterinarian.

